V-Go® Trial Form

Note: All fields below are required.

n	9	4	0	•
	a	L	C	=

Customer Details					
Customer Full Name:		Address:			
Date of Birth:		Street Address:	Suburb:		
Phone:					
Email:		State:	Postcode:		
I agree to the terms and conditions on this V-Go (Valeritas) order form. I agree that, before I use the product, I will review all the training material provided. If I need further training, I will contact my healthcare professional or AMSL Diabetes. I agree that I will make a follow up appointment with my healthcare professional.					
Hospital/Clinic Contact Details					
Prescribing Doctor Name:		Clinic/Delivery Address: Street Address:	Suburb:		
Diabetes Educator Name:					
Hospital/Clinic Name:					
Phone:					
Email:		State:	Postcode:		
I agree that the Diabetes Clinic/Hospital service listed above will provide the V-Go Insulin Patch Pump training.					
Healthcare Professional Signature:					
AMSL Diabetes Representative Name:					
Valeritas Order Details					
Billcode	Description		Qty		
VGO 40 TRIAL	V-Go 40 unit, 1 Month Trial Pack				
VGO 30 TRIAL	V-Go 30 unit, 1 Month Trial Pack				
VGO 20 TRIAL	V-Go 20 unit, 1 Month Trial Pack				

Please email completed form to diabetes@amsl.com.au

For more information on V-Go, please contact the AMSL Diabetes Customer Care Team on **1300 851 056**. amsIdiabetes.com.au (f) (2) (a) (2)

All V-Go users must review all training materials provided with the product. If the V-Go user requires further training on the product, they must contact their healthcare professional or contact AMSL Diabetes for further assistance. In agreement with the Privacy Act 1988 (Cth), the V-Go user's records will be maintained in accordance with the National Privacy Principles. The V-Go user assigns indemnity to AMSL Diabetes from and against all claims of whatsoever nature (to the maximum extent permitted by law) relating (whether directly or indirectly) to the V-Go System. V-Go Patch Pumps are not for resale. Important Risk Information. If regular adjustments or modifications to the basal rate of insulin are required in a 24-hour period, or if the amount of insulin used at meals requires adjustments of less than 2-Unit increments, use of the V-Go Disposable Insulin Delivery Device may result in hypoglycaemia. The following conditions may occur during insulin therapy with V-Go: hypoglycaemia (low blood glucose) or hyperglycemia (high blood glucose). Other adverse reactions associated with V-Go use include skin irritation from the adhesive pad or infections at the infusion site. V-Go should be removed before any magnetic resonance imaging (MRI) testing. After your initial order, you may order on our website www.amsldiabetes.com.au.

